

# Spin Physics in Out-of-Equilibrium Superconductors

**C. H. L. Quay<sup>1</sup>, D. Chevallier, M. Trif, C. Dutreix,  
M. Guigou, Y. Chiffaudel, C. Strunk\*, C. Bena<sup>†</sup>, M. Aprili.**

*Laboratoire de Physique de Solides (CNRS UMR 8502),  
Bâtiment 510, Université Paris-Sud 91405 Orsay, France.*

*\*Also at: Institute for Experimental and Applied Physics,  
University of Regensburg, 93040 Regensburg, Germany*

*<sup>†</sup>Also at: Institut de Physique Théorique, CEA Saclay 91190 Gif-sur-Yvette, France.  
charis.quay@u-psud.fr*

What happens to spin-polarised electrons when they enter a superconductor? Superconductors at equilibrium and at finite temperature contain both paired particles (of opposite spin) in the condensate phase as well as unpaired, spin-randomised quasiparticles. Injecting spin-polarised electrons into a superconductor thus creates both spin and charge imbalances (respectively  $Q^*$  and  $S^*$ ). These must relax when the injection stops, but not necessarily over the same time (or length) scale as spin relaxation requires spin-dependent interactions while charge relaxation does not. These different relaxation times can be probed by creating a dynamic equilibrium between continuous injection and relaxation, which leads to constant-in-time spin and charge imbalances. These scale with their respective relaxation times and with the injection current. While charge imbalances in superconductors have been studied in great detail both theoretically and experimentally, spin imbalances have not received much experimental attention despite intriguing theoretical predictions of spin-charge separation effects. These could occur e.g. if the spin relaxation time is longer than the charge relaxation time, i.e.  $Q^*$  relaxes faster than  $S^*$ . Fundamentally, spin-charge decoupling in superconductors is possible because the condensate acts as a particle reservoir.

We have demonstrated the existence of an almost-chargeless spin imbalance in a mesoscopic superconductor. Our measurements and fits to theory yield an estimate of the spin imbalance lifetime; we have been able to determine this quantity independently and more directly using frequency domain measurements. More recently, we have also demonstrated quasiparticle spin resonance in superconducting Al using two novel, on-chip microwave power meters.

## References

C. H. L. Quay, Y. Chiffaudel, C. Strunk and M. Aprili, ‘Quasiparticle Spin Resonance and Coherence in Superconducting Aluminium,’ submitted (2015). <http://xxx.lanl.gov/abs/1504.01615>

C. H. L. Quay, C. Dutreix, D. Chevallier, C. Bena and M. Aprili, ‘Frequency-Domain Measurement of the Spin Imbalance Lifetime in Superconductors,’ submitted (2014). <http://xxx.lanl.gov/abs/1408.1832>

D. Chevallier, C. Dutreix, M. Guigou, C. H. L. Quay, M. Aprili and C. Bena, ‘Frequency-dependent spin accumulation in out-of-equilibrium mesoscopic superconductors,’ submitted (2014). <http://xxx.lanl.gov/abs/1408.1833>

C. H. L. Quay, D. Chevallier, C. Bena and M. Aprili, ‘Spin-imbalance and spin charge separation in a mesoscopic superconductor,’ *Nature Physics*, 9, 84–88 (2013).